



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505T)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

89442-55

Date of Issuance:

7/6/22

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

TRIAD MTD

Name and Address of Registrant (include ZIP Code):

Prime Source, A Division of Albaugh LLC  
PO Box 2127  
Valdosta, GA 31604-2127

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

*Continues page 2*

Signature of Approving Official:

Date:

7/6/22

Mindy Ondish, Product Manager 23  
Herbicide Branch, Registration Division (7505P)

2. You are required to comply with the data requirements described in the generic data call-ins (GDCIs) identified below:
  - a. Triclopyr GDCI - 116001-1546
  - b. Mesotrione GDCI - 122990-1474

You must comply with all of the data requirements within the established deadlines. If you have questions about the GDCIs listed above, you may contact the Chemical Review Manager in the Pesticide Re-Evaluation Division: <http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. You have 18 months from the date of registration to provide these data.
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 12/18/2020

The following alternate brand names have been added to the product record:

- Treasure
- Fantastic
- Sublime

If you have any questions, please contact Curtis Hildebrandt by phone at (202)-566-2770, or via email at [hildebrandt.curtis@epa.gov](mailto:hildebrandt.curtis@epa.gov).

Enclosure

**ACCEPTED**

7/6/2022

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 89442-55

TRICLOPYR	GROUP	4	HERBICIDE
DICAMBA	GROUP	4	HERBICIDE
MESOTRIONE	GROUP	27	HERBICIDE

# TRIAD MTD

[Alternate Brand Names: Treasure, Fantastic, Sublime™]  
[HERBICIDE]

Provides Selective and Residual Control of Weeds in Ornamental Turfgrasses.

Active Ingredients:	% By Wt.
Triclopyr, butoxyethyl ester: 3,5,6-trichloro-2-pyridinyloxyacetic acid, butoxyethyl ester .....	29.50%
Dicamba, diglycolamine salt: diglycolamine salt of 3,6-dichloro-o-anisic acid .....	16.27%
Mesotrione: 2-[4-(methylsulfonyl)-2-nitrobenzoyl]-1,3-cyclohexanedione .....	5.20%
<b>Other Ingredients:</b> .....	<u>49.03%</u>
<b>Total:</b> .....	<b>100.00%</b>

Contains: 2.03 lbs triclopyr acid equivalent per gal; 1.00 lb dicamba acid equivalent per gal.; 0.50 lb mesotrione per gallon. Isomer specific by AOAC Methods

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

### FIRST AID

**IF SWALLOWED:**

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything to an unconscious person.

### HOTLINE NUMBER

Have the product container or label with you when calling poison control center (1-800-222-1222) or doctor, or going for treatment. For 24- Hour Emergency Assistance spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

[See inside label booklet for additional Precautionary Statements and complete Directions for Use.]

[See inside booklet for additional Precautionary Statements, complete Directions For Use, & Storage And Disposal.]

EPA Reg. No.: 89442-55

EPA Est. No.: \_\_\_\_\_

Net Contents: \_\_\_\_\_

**Manufactured [For][By]:**

Prime Source, a division of Albaugh, LLC  
1525 NE 36<sup>th</sup> Street  
Ankeny, IA 50021

# PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION.** Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse. This product may cause skin sensitization reactions in some people.

## PERSONAL PROTECTION EQUIPMENT (PPE)

**Mixers, Loaders, Applicators, and Other Handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of barrier laminate, butyl rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, natural rubber  $\geq 14$  mils, polyethylene, polyvinyl chloride  $\geq 14$  mils, or Viton  $\geq 14$  mils

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

## ENGINEERING CONTROL STATEMENTS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d-e)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

**Users should:**

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove and wash contaminated clothing before reuse.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

**GROUNDWATER ADVISORY:** Mesotrione is known to leach through soil into groundwater under certain conditions as a result of label use. Triclopyr has properties and characteristics associated with chemicals detected in groundwater. Mesotrione and triclopyr may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

**SURFACE WATER ADVISORY:** This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of mesotrione from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

**NON-TARGET ORGANISM ADVISORY:** This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

## PHYSICAL/CHEMICAL HAZARDS

Do not mix or allow this product to come in contact with oxidizing agents. Hazardous chemical reaction may occur.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI) of 24 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls worn over short-sleeved shirt and short pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of barrier laminate, butyl rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, natural rubber  $\geq 14$  mils, polyethylene, polyvinyl chloride  $\geq 14$  mils, or Viton  $\geq 14$  mils
- Chemical-resistant headgear for overhead exposure
- Protective eyewear

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard, 40 CFR Part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, and greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

### RESISTANCE MANAGEMENT

TRICLOPYR	GROUP	4	HERBICIDE
DICAMBA	GROUP	4	HERBICIDE
MESOTRIONE	GROUP	27	HERBICIDE

For resistance management, TRIAD MTD contains a Group 27, and Group 4 herbicides. Any weed population may contain or develop plants naturally resistant to TRIAD MTD and other Group 27, or 4 herbicides. The resistant biotypes may eventually dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

Fields should be scouted prior to application to identify the weed species present and their growth stage to determine if the intended application will be effective. Field should also be scouted after application to verify that the treatment was effective.

Contact your local sales representative or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of action for each target weed.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

Report any incidence of non-performance of this product against a particular weed species to your Albaugh representative or call 1-800-247-8013 or at [www.albaughLLC.com](http://www.albaughLLC.com). If resistance is suspected, treat weed escapes with an herbicide having a different mode of action and/or use non-chemical means to remove escapes, as practical, with the goal of preventing further seed production.

To delay herbicide resistance, take one or more of the following steps:

- **Diversified approach.** To the extent possible, use a diversified approach towards weed management. Whenever possible, incorporate multiple weed-control practices.
- **Know your weeds.** Identify weeds present by scouting and understand their biology. A weed-control program should consider all of the weeds present.

- **Rotate mechanisms of action.** Difficult to control weeds may require applications of herbicides with differing mechanisms of action.
- **Apply herbicide correctly.** Apply this herbicide at the correct timing and rate to control the most difficult weed in the field.

Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management directions for specific weed biotypes.

Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to these MOAs have been found in your region. **DO NOT** assume that each listed weed is being controlled by multiple mechanisms of action. Co-formulated active ingredients are intended to broaden the spectrum of weeds that are controlled. Some weeds may be controlled by only one of the active ingredients in this product.

### **PRODUCT INFORMATION**

This product is applied to weeds post-emergence to provide selective control of broadleaf weeds and certain grassy weeds in turfgrasses. This product is for use on sod farms, ornamental turf, lawns (residential, industrial, and institutional), parks, cemeteries, athletic fields, golf courses (fairways, aprons, tees, and roughs), and similar turf areas. This product can be applied to commercial and residential turfgrasses. Non-crop area use sites include golf courses, sod farms, athletic fields, parks, residential and commercial properties, cemeteries, airports, and lawns.

Post-emergent control is obtained by contact with foliage and the absorption into the plant tissue of susceptible species through contact and soil activity. Add a non-ionic surfactant when making post-emergence applications.

Before tank mixing this product with other herbicides, conduct a compatibility, safety, and efficacy test before treating larger areas. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations, and directions for use on all product labels involved in the tank mixture. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Thoroughly clean application equipment after use to avoid injury to sensitive plants.

#### **Use Restrictions:**

- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** make aerial applications.
- Retreatment Interval is 28 days.
- **DO NOT** overspray or allow spray to drift to ornamentals or flower beds and gardens. Roses and daylilies are particularly sensitive to this product.
- **DO NOT** apply more than 32 oz (0.5 lb triclopyr acid equivalent; 0.26 lb dicamba acid equivalent; 0.125 lb mesotrione) per acre per application
- **DO NOT** apply more than 128 oz (2.05 lb triclopyr acid equivalent; 1.02 lb dicamba acid equivalent; 0.5 lb mesotrione) per acre per year.
- **DO NOT** apply more than 4 applications per acre per year when using reduced application rates.
- **DO NOT** plant any crop other than turfgrass for 18 months post-application of this product.
- **DO NOT** apply organophosphate or carbamate insecticides within 7 days of applying this product.
- **DO NOT** use treated clippings to mulch trees or vegetable/flower gardens.
- **DO NOT** apply this product on Bentgrass, *Poa annua*, and kikuyugrassas injury will occur.
- **DO NOT** apply over the top of exposed roots of trees and ornamentals.
- **DO NOT** exceed specified dosages for any area.
- **DO NOT** apply to newly seeded grasses until well established.
- **DO NOT** use on golf course putting greens; maintain a minimum of a 5-foot buffer between putting greens and treated areas.
- **DO NOT** use this product on or near desirable plants including within the drip line of desirable trees and shrubs, since injury may result
- **DO NOT** apply to ditches currently being used to transport irrigation water.
- **DO NOT** apply where runoff or irrigation water may flow onto agricultural land as injury to crops may result.
- **DO NOT** apply to open water (such as lakes, reservoirs, rivers, streams, creeks, salt water bays, or estuaries).
- **DO NOT** apply this product directly to, or otherwise permit it to come into contact with or permit spray mists containing this product to drift onto cotton, grapes, tobacco, vegetable crops, flowers, fruit or orchard trees, shrubs, or other desirable broadleaf plants.
- This product is persistent and may be present in treated plant materials for over 30 days after application. **DO NOT** sell or transport treated plant materials or manure from animals that have grazed on treated plant materials off-site for compost distribution or for use as animal bedding/feed for 30 days after application.
- Animals that have been fed triclopyr treated forage must be fed forage free of triclopyr for a least 3 days before movement to an area where manure may be collected, or sensitive crops are grown.

- **DO NOT** use on sod farms in Arizona.

**PRECAUTIONS:**

- The combination of spray contact with impervious surfaces, such as roads and rocks, and increasing ambient air temperatures, may result in an increase in the volatility potential for this herbicide, increasing a risk for off-target injury to sensitive crops such as grapes and tomatoes.
- Avoid broadcast applications when air temperature exceeds 85°F. When using small, spot treatment applications in temperature over 85°F, turf injury may occur.

**MANDATORY SPRAY DRIFT MANAGEMENT**

**DO NOT** apply via air.

**Ground Boom Applications**

- **DO NOT** release spray at a height greater than 3 feet above the ground or crop canopy.
- Applicators are required to select the nozzle and pressure that deliver a medium or coarser droplet size in accordance with the American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- **DO NOT** apply during temperature inversions.

**Boom-less Ground Sprayer Applications:**

- Applicators are required to select the nozzle and pressure that deliver a medium or coarser droplet size in accordance with the American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- **DO NOT** apply during temperature inversions.

**SPRAY DRIFT ADVISORIES**

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

**IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

**Controlling Droplet Size – Ground Boom**

**Volume** – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.

**Pressure** – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.

**Spray Nozzle** – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

**BOOM HEIGHT**

**Ground Boom:** For ground equipment, the boom should remain level with the crop and have minimal bounce.

**SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

**TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

**TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

**Boomless Ground Applications**

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

**Handheld Technology Applications**  
 Take precautions to minimize spray drift.

## APPLICATION INSTRUCTIONS

**New Seedlings/New Lawn Establishment:**

Turfgrass Species	Timing	Application Rate	Comments
Kentucky bluegrass ( <i>Poa pratensis</i> ) [Buffalograss ( <i>Buchloe dactyloides</i> )] Tall fescue ( <i>Festuca arundinacea</i> ) Perennial ryegrass ( <i>Lolium perenne</i> ) [Zoysia ( <i>Zoysia spp.</i> )**] [Bahigrass ( <i>Paspalum notatum</i> )**]	Before or At Seeding from Early Spring through Fall	32 fl oz in at least 10 gallons of water per acre  (0.5 lb triclopyr ae 0.26 lb dicamba ae 0.13 lb mesotrione ai)	This product can be effectively used on grass seed blends that contain <20% by weight hard/fine fescue. For optimal control, use a NIS surfactant and apply to young, actively growing weeds prior to or at grass seeding.
Fine fescue (creeping red, chewings and hard) ( <i>Festuca spp.</i> )**	Before Seeding		
<p><b>PRECAUTION:</b>            Application to fine fescue can reduce grass density.            **Some slight visible signs of injury/discoloration may occur</p> <p><b>RESTRICTIONS:</b>  <b>DO NOT</b> spray on newly germinated turfgrass. Delay treatment until grass has been mowed 2-4 times and/or 4 weeks after emergence (whichever is longer)</p>			

**Post-Emergence Application:**

Turfgrass Species	Timing	Application Rate	Comments
Kentucky bluegrass ( <i>Poa pratensis</i> ) [Buffalograss ( <i>Buchloe dactyloides</i> )] Tall fescue ( <i>Festuca arundinacea</i> ) Perennial ryegrass ( <i>Lolium perenne</i> ) Fine fescue (creeping red, chewings and hard) ( <i>Festuca spp.</i> )** [St. Augustinegrass (grown for sod) ( <i>Stenotaphrum secundatum</i> )**] [Zoysia** ( <i>Zoysia spp.</i> )] [Centipedegrass ( <i>Eremochloa ophiuroides</i> )**] [Bahigrass ( <i>Paspalum notatum</i> )**] [Seashore Paspalum ( <i>Paspalum vagintum</i> )**]	Early Spring through Fall	32 fl oz in at least 10 gallons of water per acre  (0.5 lb triclopyr ae 0.26 lb dicamba ae 0.13 lb mesotrione ai)	For optimal control, use a NIS surfactant during application and apply to young, actively growing weeds.
<p><b>RESTRICTIONS:</b>  <b>DO NOT</b> apply this product when temperatures are above 90°F or turfgrass could result in injury.            [When applying to St. Augustinegrass (Sod uses only) and Centipedegrass, apply to established turf <b>ONLY</b>.]</p>			

[\*\*Some slight visible signs of injury/discoloration may occur]

**Dormant Bermudagrass Application (for Control of Winter Weeds):**

Turfgrass Species	Timing	Application Rate	Comments
Bermudagrass spp.	Applications above 50°F when weeds are actively growing	32 fl oz in at least 10 gallons of water per acre (0.5 lb triclopyr ae; 0.26 lb dicamba ae; 0.13 lb mesotrione ai)	Make a repeat application after 28 days.
<p><b>PRECAUTIONS:</b>            Application of this product at green-up turf will cause bermudagrass injury.</p> <p><b>RESTRICTION:</b>            Apply to fully dormant bermudagrass <b>ONLY</b>.</p>			

**Bentgrass (*Agrostis spp.*) Control:**

Weed Species	Timing	Application Rate	Comments
Bentgrass ( <i>Agrostis spp.</i> )	Applications above 50°F when weeds are actively growing	32 fl oz in at least 10 gallons of water per acre (0.5 lb triclopyr ae; 0.26 lb dicamba ae; 0.13 lb mesotrione ai)	For optimal control use a NIS surfactant. Repeat application can be made after 28 days.

**Spot Treatment Application (Sprayer):**

Spray Mix	Timing	Spray Mix Application Rate	Rate of TRIAD MTD	Rate of NIS adjuvant
-----------	--------	----------------------------	-------------------	----------------------

			<b>(per 1,000 sq. ft.)</b>	
1 gallon	Applications above 50°F when weeds are actively growing	1 gallon per 1,000 sq. ft.	0.75 fl. oz. (0.012 lb triclopyr ae; 0.006 lb dicamba ae; 0.003 lb mesotrione ai)	3 teaspoons
<b>RESTRICTION:</b> <b>DO NOT</b> apply more than 128 oz (2.03 lb triclopyr acid equivalent; 1.00 lb dicamba acid equivalent; 0.50 lb mesotrione) per acre per year.				

### Broadleaf Weeds Controlled

Alder	Coffeebean	Hoary cress	Poorjoe	Tall verbane
Amaranth	Coffeeweed	Hoary plantain	Poverty weed	Tansy mustard
Amaranth, Slender	Common chickweed	Hoary vervain	Prickly lettuce	Tansy ragwort
Amaranth, Palmer	Common groundsel	Honeysuckle	Prickly sida	Tanweed
Annual yellow sweet clover	Common mullein	Hop clover	Primrose, cutleaf evening	Thistle
Artichoke	Copperleaf	Horsenettle	Prostrate pigweed	Tick trefoil
Asiatic dayflower	Corn chamomile	Horsetail	Prostrate spurge	Toadflax
Aster spp.	Corn speedwell	Horseweed	Prostrate vervain	Trailing crown vetch
Bedstraw spp.	Cornflower	Indiana mallow	Puncture vine	Trumpercreeper
Bindweed	Creeping jenny	Ironweed	Purple cudweed	Tumble mustard
Bird vetch	Crimson clover	Jewelweed	Purple dead nettle	Tumbleweed
Birdsfoot trefoil	Croton	Jimsonweed	Purslane	Velvetleaf
Bitter sneezeweed	Cudweed	Knawel	Ragweed, common	Venice mallow
Bitter wintercress	Curly Indigo	Knotweed	Red clover	Venus lookingglass, common
Bittersweet nightshade	Dandelion	Knotweed, Giant	Redroot pigweed	Verbena
Bitterweed	Dead nettle	Knotweed, Prostrate	Redsorrel	Veronica
Black-eyed Susan	Dichondra	Knotweed, Tufted	Redstem filaree	Vervain
Black medic	Dock	Kochia	Rough cinquefoil	Vetch
Black mustard	Dock, Broadleaf	Lambsquarter	Rough fleabane	Virginia buttonweed
Black-seed plantain	Dock, Curly	Lawn burweed	Roundleaf marigold	Virginia creeper
Blessed thistle	Didder	Lespedeza	Roundleaf spurge	Virginia dwarf dandelion
Blood flower	Dogbane	Little starwort	Rush	Virginia pepperweed
Blue lettuce	Dogfennel	Locoweed	Russian pigweed	Wandering cudweed
Blue vervain	Dollar weed	Longstalked phyllanthus	Russian thistle	Wavyleaf bullthistle
Boxelder	Doveweed	Lupine	St. Johnswort	Western clematis
Bracted plantain	Eclipta	Mallow	Scarlet pimpernel	Western salsify
Brassbuttons	Elderberry	Marestail	Scotch thistle	White clover
Bristly mallow	English daisy	Marshelder	Sheep sorrel	Wild aster
Bristly oxtongue	Faceless	Matchweed	Shepherd's purse	Wild buckwheat
Broadleaf dock	Fall dandelion	Mexicanweed	Shiny cudweed	Wild carrot
Broadleaf plantain	False dandelion	Milk vetch	Slender plantain	Wild four-o'clock
Broomweed	False flax	Milkweed bloodflower	Smallflower buttercup	Wild garlic
Buckhorn	False sunflower	Morningglory	Smallflower galinsoga	Wild geranium
Buckhorn plantain	Fiddleneck	Mouseear chickweed	Smartweed	Wild lettuce
Bulbous buttercup	Field bindweed	Mugwort	Smooth dock	Wild marigold
Bull nettle	Field madder	Musk thistle	Smooth pigweed	Wild mustard
Bull thistle	Field pansy	Mustard	Sneezeweed	Wild onion
Bur ragweed	Field violet	Narrowleaf cudweed	Sorrel spp.	Wild parsnip
Burcumber	Fleabane (daisy)	Narrowleaf plantain	Southern wild rose	Wild radish
Burdock	Flixweed	Narrowleaf vetch	Sowthistle	Wild rape
Burning nettle	Florida betony	Nettle	Spanish needle	Wild strawberry
Burnweed, American	Florida pusley	Nightshade	Spatterdock	Wild sweet potato
Burweed	Frenchweed	Nimblewill	Speedwell	Wild vetch
Bushy aster	Galinsoga	Orange hawkweed	Spiderwort	Wild violet
Buttercup	Garlic mustard	Oxalis	Spiny amaranth	Willow
Canada thistle	Goathead	Oxeye daisy	Spiny cocklebur	Wintercress
Carolina false dandelion	Goatsbeard	Paleseed Plantain	Spiny sowthistle	Witchweed
Carolina geranium	Goldenrod	Parsley-piert	Spotted cats ear	Woodsorrel
Carpweed	Green foxtail	Parsnip	Spotted knapweed	Woolly croton
Catchweed bedstraw	Ground ivy	Pearlwort	Spotted spurge	Woolly morningglory
Catsear	Gumweed	Pennygrass	Spurge	Woolly plantain
Catnip	Hairy bittercress	Pennywort	Spurweed	Wormseed
Chamberbitter	Hairy beggarticks	Peppergrass	Sticky chickweed	Yarrow
Chamise	Hairy buttercup	Pepperweed	Stinging nettle	Yellow flower pepperweed
Chickweed	Hairy fleabane	Pigweed	Stinkweed	Yellow foxtail
Chicory	Hawkweed		Stitchwort	

Cinquefoil	Heal-all	Pineywoods bedstraw	Strawberry clover	Yellow nutsedge
Clover	Heartleaf drymary	Plains coreopsis	Sumac	Yellow rocket
Cockle, corn	Hedge bindweed	Plantain	Sunflower	Yellow thistle
Cockle, cow	Hedge mustard	Poison hemlock	Sweet clover	Yellow Toadflax
Cockle, white	Hemp	Poison Ivy	Swinecress	Yellow woodsorrel
Cocklebur	Henbit	Poison oak	Tall nettle	Yellowtop
		Pokeweed		

### Grassy Weeds Controlled

Common Name		Comments
Annual Bluegrass	Foxtail, Yellow	Apply to annual grasses at less than 4-tiller stage. A sequential application may be needed.
Barnyardgrass	Goosegrass*	
Bermudagrass*	Nimblewill	* May only provide suppression.
Crabgrass, Large	Signalgrass, Broadleaf	
Crabgrass, Smooth	Tufted Lovegrass	
Creeping Bentgrass	Windmillgrass	

#### PRECAUTIONS:

More mature grasses will be more difficult to control and may require a second application.

Mature, drought dressed grassy weeds (see list below) will be more difficult to control so adequate soil moisture is preferred.

Adverse or extreme environmental conditions such as poor soil conditions, high temperatures, drought and cultural conditions may affect the performance of this product.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Keep container tightly closed when not in use. Keep away from heat and flame. Do not store near seed, fertilizers, or foodstuffs. This product can be stored at temperatures as low as minus 20°F. Keep away from heat and flame.

**Pesticide Disposal:** Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Open dumping is prohibited.

**Container Handling ≤ 5 Gallons:** Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration, or by other procedures allowed by state and local authorities.

**Container Handling > 5 Gallons:** Refillable container. Refill this container with pesticide only. Do not use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration, or by other procedures allowed by state and local authorities.

**Container Handling [> 5 Gallons]:** Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

### **CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Prime Source, a division of Albaugh, LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Prime Source, a division of Albaugh LLC and Seller harmless for any claims relating to such factors.

Prime Source, a division of Albaugh, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Prime Source, a division of Albaugh, LLC, and to the extent consistent with applicable law, Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, PRIME SOURCE, A DIVISION OF ALBAUGH, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent consistent with applicable law, neither Prime Source, a division of Albaugh, LLC nor Seller shall be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF PRIME SOURCE, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF PRIME SOURCE, A DIVISION OF ALBAUGH LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Prime Source, a division of Albaugh, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Prime Source, a division of Albaugh, LLC.

063022

## LABEL HISTORY

(Not included in final printed label)

File name	Version Mark	Comment
08944-000##.20201118.DRAFT	111820	Section 3- draft label
089442-000LL.20220406.DRAFT	040622	(e) Label Revisions
089442-000LL.20220608.DRAFT	060822	(e) Label Revisions
089442-000LL.20220629.DRAFT	062922	(e) Label Revisions
089442-000LL.20220630.DRAFT	063022	(e) Label Revisions